arXiv:0801.0687v1 [math.SP] 4 Jan 2008

Borg type uniqueness Theorems for periodic Jacobi operators with matrix valued coefficients

Evgeny Korotyaev [*] Anton Kutsenko [†]

February 2, 2008

Abstract

We give a simple proof of Borg type uniqueness Theorems for periodic Jacobi operators with matrix valued coefficients.

1 Introduction

Consider a self-adjoint matrix-valued Jacobi operator J acting on $\ell^2(\mathbb{Z})^m$ and given by

$$(Jy)_n = a_n y_{n+1} + b_n y_n + a_{n-1} y_{n-1}, \quad n \in \mathbb{Z}, \quad y_n \in \mathbb{C}^m, y = (y_n)_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})^m, m \geqslant 1, \tag{1.1}$$

where $a_n > 0, b_n = b_n^*$, $n \in \mathbb{Z}$ are p-periodic sequences of the complex $m \times m$ matrices. It is well known that the spectrum $\sigma(J)$ of J is absolutely continuous and consists of non-degenerated intervals $[\lambda_{n-1}^+, \lambda_n^-], \lambda_{n-1}^+ < \lambda_n^- \leqslant \lambda_n^+, n = 1, ..., N < \infty$. These intervals are separated by the gaps $\gamma_n = (\lambda_n^-, \lambda_n^+), n = 1, .., N-1$ with the length > 0. Introduce the fundamental $m \times m$ matrix-valued solutions $\varphi = (\varphi_n(z))_{n\in\mathbb{Z}}, \vartheta = (\vartheta_n(z))_{n\in\mathbb{Z}}$ of the equation

$$a_n y_{n+1} + b_n y_n + a_{n-1}^* y_{n-1} = z y_n, \qquad \varphi_0 \equiv \vartheta_1 \equiv 0,\ \varphi_1 \equiv \vartheta_0 \equiv I_m, \quad (z,n) \in \mathbb{C} \times \mathbb{Z}, \tag{1.2}$$

where I_m is the identity $m \times m$ matrix. We define the monodromy $2m \times 2m$ matrix $\mathscr{M}_p$ by

$$\mathscr{M}_p(z) = \begin{pmatrix} \vartheta_p(z) & \varphi_p(z) \\ \vartheta_{p+1}(z) & \varphi_{p+1}(z) \end{pmatrix}, \quad z \in \mathbb{C}. \tag{1.3}$$

Let $\tau_1(z), .., \tau_{2m}(z)$ be eigenvalues of $\mathscr{M}_p(z)$. Recall that $\sigma(J) = \bigcup_{j=1}^{2m} \{z \in \mathbb{C} : |\tau_j(z)| = 1\}$, see [KKu]. Let J^0 be the unperturbed Jacobi matrix with $a_n^0 = I_m$, $b_n^0 = 0$. Note, that $\sigma(J^0) = \sigma_{ac}(J^0) = [-2, 2]$. Let $\mathbb{N}_s = \{1, .., s\}$ and let $c = \det \prod_{n=1}^p a_n$. We formulate first Borg type uniqueness theorem

---

[*] Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

[†] Department of Mathematics of Sankt-Petersburg State University, Russia e-mail: kucenkoa@rambler.u

Theorem 1.1. *Let* $c = 1$ *and let* $\sigma(J) = [\lambda_0^+, \lambda_N^-] = \{each\ |\tau_j(z)| = 1,\ (z,j) \in [\lambda_0^+, \lambda_N^-] \times \mathbb{N}_{2m}\}$*, where* $\lambda_0^+ = -\lambda_N^-$*. Then* $J = J^0$*.*

This theorem was proved in [CGR]. We give a simple proof based on the trace formula and the properties of the Chebyshev polynomials proved in Lemma 2.3. There is an enormous literature on inverse spectral problems for scalar (i.e., $m = 1$) periodic Jacobi matrices (see [BGGK], [K],[KKu1], [KKu2], [vM], [T], book [T] and references therein), but very little for matrix-valued periodic Jacobi operators (see [CGR] and references therein). Note that the complete solution of inverse problem for finite matrix-valued Jacobi operators was obtained recently [BCK].

The operator J is unitarely equivalent to the operator $\mathscr{J} = \int_{[0,2\pi)}^{\oplus} K_p(e^{ix})\frac{dx}{2\pi}$ acting in $\int_{[0,2\pi)}^{\oplus} \mathscr{H}\frac{dx}{2\pi}$, where $\mathscr{H} = \mathbb{C}^p$ and $pm \times mp$ matrix $K_p(\tau)$ is given by

$$K_p(\tau) = \begin{pmatrix} b_1 & a_1 & 0 & ... & \tau^{-1}a_p \\ a_1 & b_2 & a_2 & ... & 0 \\ 0 & a_2 & b_3 & ... & 0 \\ ... & ... & ... & ... & ... \\ \tau a_p & 0 & ... & a_{p-1} & b_p \end{pmatrix}, \quad \tau \in \mathbb{S}^1 = \{\tau : |\tau| = 1\}, \tag{1.4}$$

see [KKu]. Let $(\lambda_n(\tau))_{n=1}^{pm}$ be eigenvalues of $K_p(\tau)$. We formulate our main result.

Theorem 1.2. *Let* $a_n > 0, b_n = b_n^*$ *be real* $m \times m$ *matrices for all* $n \in \mathbb{Z}$*.*
i) Let $c = 1$*. Then* $\sum_{n=1}^{mp} \lambda_n^2(\tau) = 2pm$ *for some* $\tau \in \mathbb{S}^1$ *iff* $J = J^0$*.*
ii) Let $c = 1$ *and let* $\varkappa_1 \in \mathbb{R}$*. Then eigenvalues* $\lambda_s(e^{i\varkappa_1}) = 2\cos\frac{\varkappa_1 + 2\pi(s-1)}{p}$ *for all* $s \in \mathbb{N}_{mp}$ *iff* $J = J^0$*.*
iii) Let $\varkappa_1, \varkappa_2 \in \mathbb{R}$ *and* $\cos\varkappa_1 \neq \cos\varkappa_2$*. Let eigenvalues* $\lambda_s(e^{i\varkappa_1}) = 2\cos\frac{\varkappa_1 + 2\pi(s-1)}{p}$ *for all* $s \in \mathbb{N}_{mp}$ *and* $\lambda_n(e^{i\varkappa_2}) = 2\cos\frac{\varkappa_2 + 2\pi n_1}{p}, n \in \mathbb{N}_m$*, for some* $n_1 \in \mathbb{N}_p$*, and if* $e^{i2\varkappa_2} = 1$*, then additional eigenvalues* $\lambda_{n+m}(e^{i\varkappa_2}) = \lambda_1(e^{i\varkappa_2}) \neq \pm 2$*,* $n \in \mathbb{N}_m$*. It is possible iff* $J = J^0$*.*

Remark. 1) Note that the condition $\cos\frac{\varkappa_2 + 2\pi n_1}{p} \neq \pm 1$ in iii) is associated with the unperturbed operator J^0, where the endpoints of the spectrum $\sigma(J^0) = [-2, 2]$ have the multiplicity m, as the zeros of the determinant $D_p(z, \pm 1)$. Each point from $(-2, 2)$ has the multiplicity $2m$ as the zero of the determinant $D_p(z, \tau), \tau \neq \pm 1$.
2) Consider the case $m = 2$ and $\varkappa_1 = 0$, $\varkappa_2 = \pi$. Let periodic eigenvalues $\lambda_s(1) = 2\cos\frac{\varkappa_1 + 2\pi(s-1)}{p}$ for all $s \in \mathbb{N}_{2p}$ and let anti-periodic eigenvalues $\lambda_s(-1) = 2\cos\frac{\pi(2n+1)}{p} \neq \pm 2, s \in \mathbb{N}_4$ for some $n \in \mathbb{N}_p$. Then we deduce that $J = J^0$.
3) Consider a self-adjoint matrix-valued Jacobi operator $\widetilde{J}$ acting on $\ell^2(\mathbb{Z})^m$ and given by

$$(\widetilde{J}y)_n = \widetilde{a}_n y_{n+1} + \widetilde{b}_n y_n + \widetilde{a}_{n-1}^* y_{n-1}, \quad n \in \mathbb{Z}, \quad y_n \in \mathbb{C}^m, y = (y_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z})^m, m \geqslant 1,$$

where $\widetilde{a}_n$ ($\det\widetilde{a}_n \neq 0$), $\widetilde{b}_n = \widetilde{b}_n^*$, $n \in \mathbb{Z}$ are p-periodic sequences of the complex $m \times m$ matrices. Then the operator $\widetilde{J} = UJU^*$ where J is given by (1.2) and $U = \mathrm{diag}_{n \in \mathbb{Z}}\, u_n$ and the unitary matrices u_n have the forms $u_0 = I_m$; $u_{n+1} = c_n^* u_n$, $n \geqslant 0$; $u_{n-1} = c_{n-1} u_{n+1}$, $n \leqslant 0$. Here $\widetilde{a}_n = c_n a_n, a_n > 0$ and c_n are the unitary matrices.

2 Proof

We need the following results from [KKu].

Lemma 2.1. *The following identities and estimate are fulfilled:*

$$\sum_{n=1}^{mp}\lambda_{n,p}(\tau)=\sum_{n=1}^{p}\operatorname{Tr}b_n,\qquad \sum_{n=1}^{mp}\lambda_{n,p}^2(\tau)=\sum_{n=1}^{p}\operatorname{Tr}(b_n^2+2a_n^2),\qquad \textit{all}\quad \tau\in\mathbb{C}\setminus\{0\},\tag{2.1}$$

$$\sum_{n=1}^{pm}\lambda_{n,p}^2(\tau)\geqslant 2pmc^{\frac{2}{pm}},\qquad \tau\in\mathbb{S}^1,\tag{2.2}$$

where the identity (2.2) *holds true iff* $J=c^{\frac{1}{pm}}J^0$, *and here* $c=\det\prod_1^p a_n$.

Recall the following identities for $D_p(z,\tau)=\det(\mathscr{M}_p(z)-\tau I_{2m})$ from [KKu]

$$D_p(z,\tau)\;=\;\prod_{j=1}^{2m}(\tau_j(z)\;-\;\tau)\;=\;c(-\tau)^m\prod_{n=1}^{mp}(z\;-\;\lambda_n(\tau))\;=\;c(-\tau)^m\det(z\;-\;K_p(\tau)),\tag{2.3}$$

where $z,\tau\in\mathbb{C},\quad \tau\neq 0$.
Proof of Theorem 1.1 Consider the first case $\sum_{n=1}^p\operatorname{Tr}b_n=0$. Consider the p-periodic operator J as pk-periodic operator for some $(k,\tau)\in\mathbb{N}\times\mathbb{S}^1$. Let the eigenvalues of $K_{pk}(\tau)$ be given by

$$\lambda_-(\tau)\equiv\lambda_{1,pk}(\tau)\leqslant\lambda_{2,pk}(\tau)\leqslant..\leqslant\lambda_+(\tau)\equiv\lambda_{pmk,pk}(\tau).\tag{2.4}$$

The eigenvalues $\lambda_\pm(\tau)$ belong to $\sigma(J)$, then we have that $[\lambda_-(\tau),\lambda_+(\tau)]\subset\sigma(J)$. Using $|\tau_j(z)|=1$ for all $(z,j)\in[\lambda_-(\tau),\lambda_+(\tau)]\times\mathbb{N}_{2m}$, and (2.3), we obtain

$$|D_{pk}(z,\tau)|=\left|\prod_{j=1}^{2m}(\tau-\tau_j^k(z))\right|\leqslant\prod_{j=1}^{2m}(|\tau|+|\tau_j^k(z)|)\leqslant 2^m,\quad \textit{all}\quad z\in[\lambda_-(\tau),\lambda_+(\tau)].$$

We have $\det\prod_{n=1}^{pk}a_n=c^k=1$, since a sequence a_n is p-periodic. Then (2.3) gives

$$|D_{pk}(z,\tau)|=\left|\prod_{n=1}^{pmk}(z-\lambda_{n,pk}(\tau))\right|\leqslant 2^m,\quad \textit{all}\quad z\in[\lambda_-(\tau),\lambda_+(\tau)].\tag{2.5}$$

A sequence b_n is p-periodic, then

$$\sum_{n=1}^{pmk}\lambda_{n,pk}(\tau)=\sum_{n=1}^{pk}\operatorname{Tr}b_n=k\sum_{n=1}^{p}\operatorname{Tr}b_n=0.\tag{2.6}$$

The relations (2.4), (2.5), (2.6) and Lemma 2.3 give that

$$(\lambda_{n,pk})_{n=1}^{mpk}\in\mathcal{P}_{mpk}(2^m),\qquad \sum_{n=1}^{pmk}\lambda_{n,pk}^2(\tau)\leqslant 2pmkC_k,\qquad C_k=2^{\frac{2(m-1)}{pmk}}.\tag{2.7}$$

Then using (2.1) and $a_n = a_{n+p}, b_n = b_{n+p}$ for all $n \in \mathbb{Z}$, we obtain

$$\sum_{n=1}^{pmk} \lambda_{n,pk}^2(\tau) = \sum_{n=1}^{pk} \mathrm{Tr}(b_n^2 + a_n^2) = k \sum_{n=1}^{p} \mathrm{Tr}(b_n^2 + a_n^2) = k \sum_{n=1}^{pm} \lambda_{n,p}^2 \leqslant k(2pm)C_k,$$

which yields $\sum_{n=1}^{p} \lambda_{n,p}^2 \leqslant 2pm$, since we take $k \to +\infty$. Thus Lemma 2.1 implies $J = J^0$.

Consider the second case $\beta = \sum_{n=1}^{p} \mathrm{Tr}\, b_n$. Then a new operator $J^1 = (J - \beta)$ satisfies $\sum_{n=1}^{p} \mathrm{Tr}(b_n - \beta I_m) = 0$. Thus due to the first case we deduce that

$$\sigma(J) = \sigma(J^1) = [-2, 2] = [-x - \beta, x - \beta], \quad a_n = I_m, \quad b_n - \beta I_m = b_n^1 = 0$$

for $x = \lambda_n^-$, which yields $x = 2$, $\beta = 0$ and the theorem has been proved. ■

Lemma 2.2. *i) Let* λ *be an eigenvalue of* $K_p(\tau)$ *and have multiplicity* m *for some* $\tau \in \mathbb{S}^1 \setminus \{-1, 1\}$. *Then the multiplaers have the form* $\tau_j(\lambda) = \tau, \tau_{j+m}(\lambda) = \tau^{-1}$ *for all* $j \in \mathbb{N}_m$. *ii) Let* λ *be an eigenvalue of* $K_p(\tau)$ *and have multiplicity* $2m$ *for some* $\tau \in \{-1, 1\}$. *Then each* $\tau_j(\lambda) = \tau$, $j \in \mathbb{N}_{2m}$.

Proof. i) The matrix $K_p(\tau)$ is self-adjoint. Let $K_p(\tau) f^k = \lambda f^k$ for some orthogonal eigenvectors $f^k = (f_n^k)_{n=1}^{mp}$, $k \in \mathbb{N}_m$. If $f_0^k = \tau^{-1} f_p^k$, then the definition of the matrix $\mathscr{M}_p$ gives $\mathscr{M}_p(\lambda)(f_0^k, f_1^k)^\top = \tau(f_0^k, f_1^k)^\top$. Note that $\widetilde{f}^k = (f_0^k, f_1^k)^\top$ define other components of vector f^k, since $L(\tau)$ has special form, see (1.4). Then the vectors $\widetilde{f}^k$, $k \in \mathbb{N}_m$ are linearly independent vectors, since f^k are linearly independent vectors. Then τ has multiplicity at least m. The matrix $\mathscr{M}_p$ is symplectic, then τ^{-1} is eigenvalue of $\mathscr{M}_p$ and τ^{-1} has multiplicity at least m. We obtain first statement, since $\mathscr{M}_p$ is $2m \times 2m$ matrix. The proof of ii) is similar. ■

Proof of Theorem 1.2. The statement i) follows from Lemma 2.1.

ii) **Sufficiency**. Recall that if $m = 1$, then $\det(M_p^0 - \tau I_2) = \tau(\tau + \tau^{-1} - 2\mathscr{T}_p(z/2))$, where $\mathscr{T}_p(z) = \cos(p \arccos z)$ is the Chebyshev polynomial. Moreover, zeros of the polynomial $\mathscr{T}_p(z) - \cos \varkappa_1$ are given by $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$, $s \in \mathbb{N}_p$.

Thus if $J = J^0, m \geqslant 2$, then corresponding monodromy operators M_p satisfies $\det(M_p - \tau I_{2m}) = \tau^m \prod_{j=1}^{m} (\tau + \tau^{-1} - 2\mathscr{T}_p(z/2))$, which yields at $\tau = e^{i\varkappa_1}$

$$det(M_p - e^{i\varkappa_1} I_{2m}) = \tau = e^{im\varkappa_1} 2^m (\cos \varkappa_1 - \mathscr{T}_p(z/2))^m = e^{im\varkappa_1} 2^m, \tag{2.8}$$

This implies $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$, $s \in \mathbb{N}_{mp}$.

Necessity. Let $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$, $s \in \mathbb{N}_{mp}$. Then the direct calculation implies $\sum_{n=1}^{mp} \lambda_n^2(e^{i\varkappa_1}) = 2pm$, and the statement i) gives $J = J^0$.

iii) **The sufficiency** is proved similar to the case ii).

Necessity. We consider only the case $e^{2i\varkappa_2} \neq 1$, the proof of the case $e^{2i\varkappa_2} \neq 1$ is similar. Let $\lambda_s(e^{i\varkappa_1}) = 2 \cos \frac{\varkappa_1 + 2\pi(s-1)}{p}$, $s \in \mathbb{N}_{mp}$. Using (2.3), (2.8), we obtain

$$\prod_{j=1}^{2m} (e^{i\varkappa_1} - \tau_j(z)) = c(-1)^m e^{im\varkappa_1} \prod_{n=1}^{mp} (z - \lambda_n(e^{i\varkappa_1})) = c(-2)^m e^{im\varkappa_1} (\mathscr{T}_p(z/2) - \cos \varkappa_1)^m, \tag{2.9}$$

where $c = \det \prod_{n=1}^{p} a_n$. The eigenvalue $\lambda_1(e^{i\varkappa_2})$ has the multiplicity m, then using Lemma 2.2 and substituting $z = \lambda_1(e^{i\varkappa_2}) = 2\cos\frac{\varkappa_2 + 2\pi n_1}{p}$ and two multipliers $e^{\pm i\varkappa_2}$ (given by Lemma 2.2) into (2.9), we obtain

$$(e^{i\varkappa_1} - e^{i\varkappa_2})^m (e^{i\varkappa_1} - e^{-i\varkappa_2})^m = c2^m e^{im\varkappa_1} (\cos\varkappa_2 - \cos\varkappa_1)^m,$$

which yields $c = 1$. Then the statement ii) gives $J = J^0$. ■

Lemma 2.3. *For any $r \geqslant 0$, $s \geqslant 2$ the following identity holds true*

$$\sup_{x \in \mathcal{P}_s(r)} \sum_1^s x_n^2 = 2s \left(\frac{r}{2}\right)^{\frac{2}{s}},$$

$$\mathcal{P}_s(r) = \left\{ (x_n)_1^s : \ x_1 \leqslant .. \leqslant x_s, \ \sum_{n=1}^{s} x_n = 0, \ \left| \prod_{n=1}^{s} (z - x_n) \right| \leqslant r, \quad all \ z \in [x_1, x_s] \right\} \subset \mathbb{R}^s.$$

Proof. In the proof we use arguments from [Ku]. Let $\|x\|^2 = \sum x_n^2, x = (x_n)_1^s \in \mathbb{R}^s$. The set $\mathcal{P}_s(r)$ is compact, then $\sup_{x \in \mathcal{P}_s(r)} \|x\|^2 = \|x^0\|^2$ for some $x^0 = (x_n^0)_1^s \in \mathcal{P}_s(r)$. Introduce a polynomial $p_0(z) = \prod_{n=1}^{s}(z - x_n^0)$. Let $x_0^0 = -\infty$, $x_{s+1}^0 = +\infty$. The polynomial p_0' has only real zeros $\widetilde{x}_n, n \in \mathbb{N}_{s-1}$. We will show that each $|p_0(\widetilde{x}_n)| = r, n \in \mathbb{N}_{s-1}$. Assume that there exist $1 \leqslant n_1 < n_2 \leqslant s$ such that

$$x_{n_1-1}^0 < x_{n_1}^0 \leqslant x_{n_2}^0 < x_{n_2+1}^0, \quad \max_{z \in [x_{n_1}, x_{n_2}]} |p_0(z)| < r. \tag{2.10}$$

Introduce a polynomial $p_\varepsilon(z) = \prod_{n=1}^{s}(z - x_n^\varepsilon)$, where a vector $x^\varepsilon = (x_n^\varepsilon)_{n=1}^s \in \mathbb{R}^s$ is given by

$$x_n^\varepsilon = x_n^0, \ n \neq n_1, \ n \neq n_2; \quad x_{n_1}^\varepsilon = x_{n_1}^0 - \varepsilon, \ x_{n_2}^\varepsilon = x_{n_2}^0 + \varepsilon, \quad \varepsilon > 0. \tag{2.11}$$

Using $x^0 \in \mathcal{P}_s(r)$ and (2.10), we obtain

$$x_1^\varepsilon \leqslant .. \leqslant x_s^\varepsilon, \quad \sum_{n=1}^{s} x_n^\varepsilon = \sum_{n=1}^{s} x_n^0 = 0, \quad \max_{z \in [x_{n_1}^\varepsilon, x_{n_2}^\varepsilon]} |p_\varepsilon(z)| \leqslant r, \tag{2.12}$$

for sufficiently small $\varepsilon > 0$. We rewrite p_ε in the form

$$p_\varepsilon(z) = p_0(z) g_\varepsilon(z), \quad g_\varepsilon(z) = \frac{(z - x_{n_1}^\varepsilon)(z - x_{n_2}^\varepsilon)}{(z - x_{n_1}^0)(z - x_{n_2}^0)} = \frac{(z - x_{n_1}^0 + \varepsilon)(z - x_{n_2}^0 - \varepsilon)}{(z - x_{n_1}^0)(z - x_{n_2}^0)}. \tag{2.13}$$

Due to $x_{n_1}^0 \leqslant x_{n_2}^0$ and $\varepsilon > 0$ we deduce that $|g_\varepsilon(z)| \leqslant 1$, $z \in \mathbb{R} \setminus [x_{n_1}^\varepsilon, x_{n_2}^\varepsilon]$. Then (2.13) yields $|p_\varepsilon(z)| \leqslant |p_0(z)|$, $z \in \mathbb{R} \setminus [x_{n_1}^\varepsilon, x_{n_2}^\varepsilon]$ and thus (2.12) gives $|p_\varepsilon(z)| \leqslant r$, $z \in [x_1^\varepsilon, x_s^\varepsilon]$, and $x^\varepsilon \in \mathcal{P}_s(r)$ for all sufficiently small $\varepsilon > 0$. There is an estimate

$$\|x^\varepsilon\|^2 = \sum_{n=1}^{s} (x_n^\varepsilon)^2 = \sum_{n=1}^{s} (x_n^0)^2 + 4\varepsilon(x_{n_2}^0 - x_{n_1}^0) + 2\varepsilon^2 > \|x^0\|^2, \tag{2.14}$$

since $x_{n_1}^0 \leqslant x_{n_2}^0$ and $\varepsilon > 0$. But (2.14) and the condition $x^\varepsilon \in \mathcal{P}_s(r)$ contradict the identiy $\sup_{x\in\mathcal{P}_s(r)} \|x\|^2 = \|x^0\|^2$. Then the assumption (2.10) is not true and each $|p_0(\widetilde{x}_n)| = r, n \in \mathbb{N}_{s-1}$. Note that only polynomials $r\mathscr{T}_s(\alpha z + \beta)$, $\alpha \in \mathbb{R} \setminus \{0\}$, $\beta \in \mathbb{R}$, (here $\mathscr{T}_s$ are the Chebyshev polynomials, i.e. $\mathscr{T}_s(\cos z) = \cos sz$) have this property, then $p_0(z) = r\mathscr{T}_s(\alpha z+\beta)$ for some $\alpha \in \mathbb{R} \setminus \{0\}$, $\beta \in \mathbb{R}$. We take α, β such that $p_0(z) = \prod_{n=1}^s (z - x_n^0)$, $\sum_{n=1}^s x_n^0 = 0$ and then $p_0(z) = r\mathscr{T}_s(zr^{-\frac{1}{s}}2^{\frac{1-s}{s}})$, since (see [AS])

$$\mathscr{T}_s(z) = \frac{1}{2}\sum_{k=0}^{[\frac{s}{2}]}(-1)^k \frac{s}{s-k} C_k^{s-k}(2z)^{s-2k} = 2^{s-1}z^s - 2^{s-3}sz^{s-2} + o(z^{s-2}) \quad as\ z \to \infty,$$

where $C_m^n = \frac{n!}{m!(n-m)!}$. Then using Viette formulas we get

$$\mathscr{T}_s(z) = 2^{s-1}\prod_{n=1}^{s}(z - z_n) = 2^{s-1}\left(z^s - \xi z^{s-1} + \frac{1}{2}\left(\xi^2 - \eta\right) z^{s-2} + o(z^{s-2})\right) \quad as\ z \to \infty,$$

where z_n, $n \in \mathbb{N}_s$ are zeroes of $\mathscr{T}_s$ and $\xi = \sum_1^s z_n$, $\eta = \sum_1^s z_n^2$. Then we get $\xi = 0$ and $\eta = \frac{s}{2}$. This gives

$$\sup_{x\in\mathcal{P}_s(r)} \|x\|^2 = \|x^0\|^2 = r^{\frac{2}{s}} 2^{\frac{2s-2}{s}} \sum_1^s z_n^2 = r^{\frac{2}{s}} 2^{\frac{2s-2}{s}} \eta = 2s\left(\frac{r}{2}\right)^{\frac{2}{s}} \blacksquare$$

Acknowledgments. Evgeny Korotyaev was partly supported by DFG project BR691/23-1. The some part of this paper was written at the Math. Institute of Humboldt Univ., Berlin; Anton Kutsenko is grateful to the Institute for the hospitality.

References

[AS] Abramowitz, M.; Stegun, A., eds. Handbook of Mathematical Functions. N.Y.: Dover Publications Inc., 1992.

[BGGK] Bättig, D.; Grebert, B.; Guillot, J.-C.; Kappeler, T. Fibration of the phase space of the periodic Toda lattice. J. Math. Pures Appl. 72 (1993), no. 6, 553–565.

[BCK] Brüning,J.; Chelkak, D.; Korotyaev, E. Inverse spectral analysis for finite matrix-valued Jacobi operators, preprint 2006.

[CG] Clark, S.; Gesztesy, F. On Weyl–Titchmarsh theory for singular finite difference Hamiltonian systems, J. Comput. Appl. Math., 171 (2004), 151–184.

[CGR] Clark, S.; Gesztesy, F.; Renger, W. Trace formulas and Borg-type theorems for matrix-valued Jacobi and Dirac finite difference operators. J. Differential Equations 219 (2005), no. 1, 144–182.

[GKM] Gesztesy, F.; Kiselev, A.; Makarov, K.,A. Uniqueness results for matrix-valued Schrödinger, Jacobi, and Dirac-type operators, Math. Nachr. 239–240 (2002), 103–145.

[K] Korotyaev, E. Gap-length mapping for periodic Jacobi matrices, Russ. J. Math. Phys. 13(2006), no.1, 64-69.

[KKr] Korotyaev, E.; Krasovsky, I. Spectral estimates for periodic Jacobi matrices, Commun. Math. Phys. 234 (2003), 517-532.

[KKu] Korotyaev, E.; Kutsenko, A. Lyapunov functions for periodic matrix-valued Jacobi operators, preprint 2007, www.arxiv.org.

[KKu1] Korotyaev, E.; Kutsenko, A. Inverse problem for the discrete 1D Schrödinger operator with small periodic potentials, Commun. Math. Phys. 261(2006), 673-692.

[KKu2] Korotyaev, E.; Kutsenko, A. Marchenko-Ostrovski mappings for periodic Jacobi matrices, will be published in Russ. J. Math. Phys.

[Ku] Kutsenko, A. Estimates of parameters for conformal mappings related to a periodic Jacobi matrix. Journal of Mathematical Sciences, 134 (2006), no. 4, 2295–2304.

[vM] van Moerbeke, P. The spectrum of Jacobi matrices. Invent. Math. 37 (1976), no. 1, 45–81.

[T] Teschl, G. Jacobi Operators and Completely Integrable Nonlinear Lattices, Mathematical Survey Monographs, vol. 72, American Mathematical Society, Providence, RI, 2000.